



**S**hared  
**H**ealthcare  
**I**ntervention to  
**E**liminate  
**L**ife-threatening  
**D**issemination of MDROs

**Hospital Protocol Post-Training Test:  
Chlorhexidine basin bed bathing with CHG liquid  
and nasal iodophor administration**

Date: \_\_\_\_\_ Title (*circle one*): CNA LVN RN OTHER: \_\_\_\_\_

Name: \_\_\_\_\_ Signature: \_\_\_\_\_

**1. What is the SHIELD Program?**

- A. A quality improvement program to reduce multi-drug resistant organisms (MDROs) like MRSA and ESBL in hospitals
- B. A research project to minimize the spread of multi-drug resistant organisms
- C. A project to evaluate novel decolonization products for FDA clearance
- D. An initiative to improve bathing techniques for all patients

**2. Who should receive the decolonization "cleansing" protocol?**

- A. MRSA-positive patients
- B. Patients on contact precautions
- C. Patients who stay longer than 2 weeks
- D. All patients over 18 years old

**3. How much water and 4% liquid CHG do you mix together when performing basin bed bath?**

- A. One 4 oz bottle of 4% CHG, one 4 oz bottle of water
- B. Two 4 oz bottles of 4% CHG, two 4 oz bottles of water
- C. One 4 oz bottle of 4% CHG, two 4 oz bottles of water

**4. Which of the following statement is FALSE when using basin bed bathing with CHG?**

- A. Do not dilute more than equal part of water to CHG
- B. Wring each wipe soaked in basin before application
- C. Soak used wipes in basin and apply again

- 5. Which of the following is TRUE when using basin bed bathing with chlorhexidine?**
- A. Apply chlorhexidine wipes to the skin and then bathe with soap and water
  - B. Bathe with soap and water then apply chlorhexidine
  - C. Bathe with chlorhexidine cloths and let air dry
- 6. Chlorhexidine liquid can be used on all of the following EXCEPT:**
- A. Lines, tubes, and drains within 6 inches of the body
  - B. Superficial decubitus ulcers (stage 1&2)
  - C. Large or deep wounds
  - D. Sutured or stapled wounds
- 7. Which dressings should be cleaned with chlorhexidine?**
- A. Gauze dressings
  - B. Semi-permeable or occlusive dressings
- 8. Do not use soap to cleanse incontinent patients because soap can inactivate CHG.**
- A. True
  - B. False
- 9. Who should receive nasal iodophor?**
- A. Nasal iodophor should be given to MRSA-positive patients only
  - B. Nasal iodophor should be given to all patients on contact precautions
  - C. Nasal iodophor should be given to all patients
- 10. Which of the following is correct when using nasal iodophor?**
- A. Apply once to each nostril
  - B. Apply twice daily to each nostril for five days
  - C. Apply only to patients known to have MRSA



**S**hared  
**H**ealthcare  
**I**ntervention to  
**E**liminate  
**L**ife-threatening  
**D**issemination of MDROs

**Hospital Protocol Post-Training Test Answer Key:  
Chlorhexidine basin bed bathing with CHG liquid  
and nasal iodophor administration**

- 1. A
- 2. B
- 3. A
- 4. C
- 5. C
- 6. C
- 7. B
- 8. A
- 9. B
- 10. B

