Prevent infections during hospital stay


BASIN BED BATHING with Chlorhexidine (CHG) Liquid



Bathe with CHG to remove germs and prevent infection

CHG works better than soap and water
CHG is a protective bath

Apply as shown below



Avoid eyes, mouth, & ear canals

BASIN BATH Instructions

- 1. Prepare one 4 oz bottle of 4% liquid CHG, a bed basin, and 6 disposable wipes (more if needed).
- 2. Dispense one 4 oz bottle of 4% CHG liquid into basin.



- Fill one 4 oz bottle with water and dispense into basin. Do not overdilute.
 Use equal part of water to CHG.
- 4. Soak wipes in basin and wring before use. Do not place back into basin after use.
- 5. **Firmly massage** skin with wipes.
- 6. Clean over semi-permeable dressings.
- 7. Clean 6 inches of lines, tubes, and drains.

REMINDERS

- Your enthusiasm helps patients understand why CHG is important
- Bathing on admission removes germs to protect the patient and hospital
- CHG works for 24 hours to kill germs
- Firmly massage CHG onto skin
- Clean **6 inches** of lines, drains, tubes
- Safe on surface wounds, rashes, burns
- Use only CHG-compatible lotions
- If barrier protection needed, apply CHG then apply barrier protection

Clean all skin areas with attention to:

- Neck
- All skin folds
- Skin around all devices (line/tube/drain)
- Wounds unless deep or large
- Armpit, groin, between fingers/toes