



Shared
Healthcare
Intervention to
Eliminate
Life-threatening
Dissemination of MDROs

**LTACH Protocol Post-Training Test:
Chlorhexidine basin bed bathing with CHG liquid
and nasal iodophor administration**

Date: _____ Title (*circle one*): CNA LVN RN OTHER: _____

Name: _____ Signature: _____

1. What is the SHIELD Program?

- A. A quality improvement program to reduce multi-drug resistant organisms (MDROs) like MRSA and ESBL in hospitals
- B. A research project to minimize the spread of multi-drug resistant organisms
- C. A project to evaluate novel decolonization products for FDA clearance
- D. An initiative to improve bathing techniques for all patients

2. Who should receive the decolonization "cleansing" protocol?

- A. MRSA-positive patients
- B. Patients on contact precautions
- C. Patients who stay longer than 2 weeks
- D. All patients over 18 years old

3. How much water and 4% liquid CHG do you mix together when performing basin bed bath?

- A. One 4 oz bottle of 4% CHG, one 4 oz bottle of water
- B. Two 4 oz bottles of 4% CHG, two 4 oz bottles of water
- C. One 4 oz bottle of 4% CHG, two 4 oz bottles of water

4. Which of the following statement is FALSE when using basin bed bathing with CHG?

- A. Do not dilute more than equal part of water to CHG
- B. Wring each wipe soaked in basin before application
- C. Soak used wipes in basin and apply again

- 5. Which of the following is TRUE when using basin bed bathing with chlorhexidine?**
- A. Apply chlorhexidine wipes to the skin and then bathe with soap and water
 - B. Bathe with soap and water then apply chlorhexidine
 - C. Bathe with chlorhexidine cloths and let air dry
- 6. Chlorhexidine liquid can be used on all of the following EXCEPT:**
- A. Lines, tubes, and drains within 6 inches of the body
 - B. Superficial decubitus ulcers (stage 1&2)
 - C. Large or deep wounds
 - D. Sutured or stapled wounds
- 7. Which dressings should be cleaned with chlorhexidine?**
- A. Gauze dressings
 - B. Semi-permeable or occlusive dressings
- 8. Do not use soap to cleanse incontinent patients because soap can inactivate CHG.**
- A. True
 - B. False
- 9. Who should receive nasal iodophor?**
- A. Nasal iodophor should be given to MRSA-positive patients only
 - B. Nasal iodophor should be given to all patients on contact precautions
 - C. Nasal iodophor should be given to all patients
- 10. Which of the following is correct when using nasal iodophor?**
- A. Apply once to each nostril
 - B. Apply twice daily to each nostril for five days
 - C. Apply only to patients known to have MRSA



Shared
Healthcare
Intervention to
Eliminate
Life-threatening
Dissemination of MDROs

**Hospital Protocol Post-Training Test Answer Key:
Chlorhexidine basin bed bathing with CHG liquid
and nasal iodophor administration**

1. A
2. D
3. A
4. C
5. C
6. C
7. B
8. A
9. C
10. B

