



Shared
Healthcare
Intervention to
Eliminate
Life-threatening
Dissemination of MDROs

**Nursing Home Protocol Post-Training Test:
Chlorhexidine basin bed bathing with CHG liquid
and nasal iodophor administration**

Date: _____ Title (*circle one*): CNA LVN RN OTHER: _____

Name: _____ Signature: _____

1. What is the SHIELD Program?

- A. A quality improvement program to reduce multi-drug resistant organisms (MDROs) like MRSA and ESBL in nursing homes
- B. A research project to minimize the spread of multi-drug resistant organisms
- C. A project to evaluate novel decolonization products for FDA clearance
- D. An initiative to improve bathing techniques for all residents

2. Who should receive the decolonization "cleansing" protocol?

- A. MRSA-positive residents
- B. Residents on contact precautions
- C. Residents who stay longer than 2 weeks
- D. All residents over 18 years old

3. How much water and 4% liquid CHG do you mix together when performing basin bed bath?

- A. 1 cup of CHG, ½ cup of water
- B. ½ cup of CHG, ½ cup of water
- C. ½ cup of CHG, 1 cup of water

4. Which of the following statement is FALSE when using basin bed bathing with CHG?

- A. Do not dilute more than equal part of water to CHG
- B. Wring each wipe soaked in basin before application
- C. Soak used wipes in basin and apply again

- 5. Which of the following is TRUE when using basin bed bathing with chlorhexidine?**
- A. Apply chlorhexidine wipes to the skin and then bathe with soap and water
 - B. Bathe with soap and water then apply chlorhexidine
 - C. Bathe with chlorhexidine cloths and let air dry
- 6. Chlorhexidine liquid can be used on all of the following EXCEPT:**
- A. Lines, tubes, and drains within 6 inches of the body
 - B. Superficial decubitus ulcers (stage 1&2)
 - C. Large or deep wounds
 - D. Sutured or stapled wounds
- 7. Which dressings should be cleaned with chlorhexidine?**
- A. Gauze dressings
 - B. Semi-permeable or occlusive dressings
- 8. Do not use soap to cleanse incontinent patients because soap can inactivate CHG.**
- A. True
 - B. False
- 9. Who should receive nasal iodophor?**
- A. Nasal iodophor should be given to MRSA-positive residents only
 - B. Nasal iodophor should be given to all residents on contact precautions
 - C. Nasal iodophor should be given to all residents
- 10. Which of the following is correct when using nasal iodophor?**
- A. Apply once to each nostril
 - B. Apply twice daily to each nostril for five days
 - C. Apply only to residents known to have MRSA



Shared
Healthcare
Intervention to
Eliminate
Life-threatening
Dissemination of MDROs

**Nursing Home Protocol Post-Training Test Answer Key:
Chlorhexidine basin bed bathing with CHG liquid
and nasal iodophor administration**

1. A
2. D
3. B
4. C
5. C
6. C
7. B
8. A
9. C
10. B